

FREE VACCINES SUPPLIED BY DHS VICTORIA AND THEIR INDICATION FOR USE

VACCINE	INDICATION FOR USE
Adult Diphtheria & Tetanus Adsorbed (ADT Booster) [®]	<ul style="list-style-type: none"> - Catch-up immunisation for school aged children only who have not had a primary course of diphtheria and tetanus or has an incomplete history of having diphtheria and tetanus. - Booster dose at 50 years of age only
Adult/Adolescent Diphtheria, Tetanus & acellular Pertussis (Boostrix) [®]	<ul style="list-style-type: none"> - Local council offer Boostrix[®] in Year 10 of secondary school. Single dose for year 10 students only - GPs are supplied a small quantity for Year 10 students only or age equivalent who missed having the vaccine at school
Bacille Calmette Geurin (BCG) [®]	<ul style="list-style-type: none"> - 1 vial = 10 doses - Only given by TB registered medical personnel - Infants and children less than 5 years of age in an "at risk" country for more than 3 months.
Diphtheria, Tetanus, Acellular Pertussis & Inactivated Polio (Infanrix IPV) [®]	<ul style="list-style-type: none"> - Given at 2, 4, 6 months and 4 years of age - 'Catch-up' to the 8th birthday
Inactivated Poliomyelitis (IPOL) [®]	<ul style="list-style-type: none"> - Given to school aged children only (aged 8 years and older) who are unimmunised or incompletely immunised for poliomyelitis - Call DHS to discuss before ordering
<i>Haemophilus influenzae</i> type b (PedvaxHIB) [®]	<ul style="list-style-type: none"> - For children up to their 5th birthday as a "catch-up" program - Call DHS to discuss before ordering
Hepatitis B adult (HBVax II Adult) [®]	<ul style="list-style-type: none"> - Local council offer Hepatitis B immunisation in Year 7 of secondary school; students receive 2 doses of adult hepatitis B vaccine, 4 to 6 months apart - GPs are supplied a small quantity for Year 7 students who did not receive it in the school program - 'Catch-up' for adolescents in secondary school who missed the Year 7 hepatitis B course & are under 16 years of age - Available for household contacts of a hepatitis B carrier and also for an Injecting drug user
Hepatitis B paediatric (HBVax II Paediatric) [®]	<ul style="list-style-type: none"> - Infants at birth - For children born after 1st May 2000 who have not received a course of hepatitis B vaccine - 'Catch-up' for adolescents in secondary school who missed the Year 7 hepatitis B course and are over 15 years of age - Available for household contacts (children) of a hepatitis B carrier and also for an Injecting drug user - Call DHS to discuss before ordering
Hib - Hepatitis B (Comvax) [®]	<ul style="list-style-type: none"> - Given at 2, 4 and 12 months of age
Human Tuberculin 100u/ml (10 ml vial)	<ul style="list-style-type: none"> - As per TB Guidelines
Human papillomavirus (Gardasil) [®]	<ul style="list-style-type: none"> - Local council provide vaccine to females in Year 7 of secondary school - Catch-up program for females in Years 8 – 12 of secondary school (phased rollout) - Catch-up program for females aged 18-26 years, Medicare card holders only - GPs are supplied a small quantity for students who did not receive vaccine through the school program - Catch-up program finishes June 2009
Influenza	<ul style="list-style-type: none"> - Given at 65 years and over - Given at 50 years and over for Aboriginal and Torres Strait Islanders (ATSI) - Given at 15 to 49 years for ATSI with underlying medical risk factors
Measles, Mumps, Rubella (Priorix) [®]	<ul style="list-style-type: none"> - Given at 12 months and 4 years of age - For women planning pregnancy and post partum with low or negative rubella antibody - 'Catch-up' for school aged children only with no documented history of having received two doses
Meningococcal C (NeisVac C) [®]	<ul style="list-style-type: none"> - Given at 12 months of age - 'Catch up' for children born on or after 1 January 2002
Childhood Pneumococcal (Prevenar) [®]	<ul style="list-style-type: none"> - Given at 2, 4, & 6 months of age - Children with underlying medical risk factors up to the 5th birthday
Pneumococcal (Pneumovax 23) [®]	<ul style="list-style-type: none"> - Given to non-indigenous aged 65 years and over - Given at 50 years and over for Aboriginal and Torres Strait Islanders (ATSI) - Given at 15 to 49 years for ATSI with underlying medical risk factors - Given to children with underlying medical risk factors at 4-5 years of age
Rotavirus (RotaTeq) [®]	<ul style="list-style-type: none"> - Given at 2, 4 and 6 months of age
Varicella (chickenpox) (Varilrix) [®]	<ul style="list-style-type: none"> - Given at 18 months of age - 'Catch up' for children born on or after 1 May 2004 aged at least 18 months or older - Local council offer Varilrix[®] in Year 7 of secondary school for children who have not had the vaccine or the disease - GPs are supplied a small quantity for Year 7 students only who did not receive it through the school program

Note: Vaccines required for travel purposes are not funded by DHS Victoria